

[illegible]

Schnee, S.C., Smrekar, S.E., Broecker, D.L., Ginn, R.R., Koala, W.M., McGill, G.E., Phillips, R.J., Saunders, R.S., Schertzer, G., Seayoum, S.W., and Stofan, E.R.: 1992, Venus Express: A mission to study the geology and atmosphere of Venus, *J. Geophys. Res.* **97**, 1319-1326.

Seayoum, S.W., McGill, D.G., Simons, M., Sedokov, S.C., Hager, B.H., and McGill, G.E.: 1990, Plains tectonics on Venus. The deformation belts of Lactoria Planitia Journal of Geophysical Research, v. 97, p. 13579-13610.

Seayoum, S.W., McGill, D.G., Simons, M., Beer, G., Broecker, D.L., Jares, D.M., and Scott, S.D.: 1989, Global distribution and characteristics of coronae and related features on Venus: Implications for origin and relation to mantle processes. Journal of Geophysical Research, v. 94, p. 13539-13552.

Shaw, R.T., Schaber, G.G., and Dawson, D.D.: 1994, The global resurfacing of Venus. Journal of Geophysical Research, v. 99, p. 10899-10910.

Simons, M., Seayoum, S.W., and McGill, D.G.: 1990, Basalts, basaltic dykes, and basalts, V.N. eds., *Venus: geology, geomorphology, and geophysics*: Tucson, University of Arizona Press, p. 82-95.

USGS: 1997, *Map Compiler*. *Venus geologic mapppers handbook*, second edition: U.S. Geological Survey Report No. 94-38, 50 pp.

Yager, G.L., Simpson, R.A., Maurer, M.J., and Hofmann, E.: 1992, Scattering properties of the Venusian surface from Magellan. *Journal of Geophysical Research*, v. 97, p. 11335-11340.

INTERIOR—GEOLOGICAL SURVEY RESTON, VA—2001

[illegible]

GENERAL GEOLOGY

Several different geologic processes have influenced the Lavitra Planitia region and have combined together to form its geologic record. Volcanism is the dominant process of crustal growth on Venus (Head and others, 1992) and production of the observed geologic units in this region. Tectonic activity has modified some of these basic crustal materials (for example, Solomon and others, 1992; Sogard and others, 1992) in a variety of modes (extension, contraction, and shear). In places deformation is so extensive, as in the case of tessera terrain, that the deformation has become part of the edification of the terrain (see Tanaka, 1994; Scott and others, 1986). Impact cratering also has locally influenced regions in the quadrangle, most notably areas dominated by Danilovs, Saskia, and Aglaiezer craters, but in general has not been as tidal process over the quadrangle as a whole. Eolian processes require a source of sedimentary deposits and thus are concentrated around impact craters and localized around fractures and scarps (for example, Greeley and others, 1992).

[illegible][illegible][illegible][illegible][illegible][illegible][illegible]

and the abnor-
malities of the
central part of
adjacent to the
St. Petersburg
of per unit, which
plants material
of the top of the
top of the top of
the ridged
formed by the